



Endoscopy Department
1509 29th Street South
Great Falls, MT 59405
www.gfclinic.com
Phone: (406) 771-3400

Welcome to Great Falls Hospital Endoscopy Department!

To make your procedure as safe and efficient as possible, please review the enclosed information. If you need to cancel or reschedule your procedure, or have questions, please call (406) 771-3400.

- A nurse will call you prior to your procedure to complete the pre-admission questionnaire.
- Please bring a current list of medications and known allergies with you on the day of procedure. A sheet is included in this packet to help you.
- Please bring a photo ID and insurance card on day of procedure.

Your procedure is scheduled for (date/time): _____ a.m / p.m

Please arrive at _____ a.m / p.m

Your procedure provider is: _____

**Location: Great Falls Hospital
3010 15th Ave South
Great Falls, Mt 59405**

OR

**Great Falls Hospital Surgery Center
1509 29th Street South
Great Falls, MT 59405**

Colonoscopy

Prior to Procedure:

- 1) Our prior authorization team will work on obtaining procedure authorization through your insurance company. It is still recommended to contact your insurance company to make sure authorization is in place. You may still be responsible for a deductible or co-pay. Plan to bring your insurance card and photo ID with you when checking in for your procedure.
- 2) If you take a blood thinner, such as **Brilinta (ticagrelor)**, **Coumadin (warfarin)**, **Effient (prasugrel)**, **Eliquis (apixaban)**, **Elmiron (pentosan polysulfate sodium)**, **Plavix (clopidogrel)**, **Pletal (cilostazol)**, **Pradaxa (dabigatran)**, **Savaysa (edoxaban)**, **Xarelto (rivaroxaban)**, or any other prescription blood thinner, please contact the provider who prescribes this medication for instructions as to when or if you should stop taking it prior to your procedure. Aspirin typically does not need to be stopped unless you have been specifically instructed to do so by your healthcare provider.
- 3) Please stop taking the following medications **5 days prior to your procedure**:
 - a. Iron containing medications including multivitamin with iron.
 - b. Fish Oil
 - c. Any type of fiber supplement, such as Benefiber, Metamucil, Citrucel, Psyllium, or FiberCon.
- 4) Phentermine – **hold 4 days prior to procedure**
- 5) If you have diabetes, your medication dose may require adjustments. Please contact your prescribing provider for instructions. This includes insulin and oral diabetic medications.
- 6) You will not be able to drive after the procedure or for the remainder of the day. You must have a driver over the age of 18 with you. If using a taxi or similar service, you must have an adult over the age of 18 accompany you. It is recommended that someone stay with you until the next morning. Do not operate machinery, go to work, or sign legal documents for the rest of the day. You should be able to return to work the next day.
- 7) Wear comfortable clothing the day of the procedure. Leave all valuables at home. Bring a case if you wear glasses.

Preparing for your Colonoscopy Procedure

Five (5) days before your procedure:

Low-residue diet:

You may eat foods such as:

- Chicken
- Fish
- Eggs
- White bread
- White rice
- Pasta
- Potatoes without skins
- Plain yogurt
- Broth
- Gelatin

Avoid:

- Nuts
- Seeds
- Popcorn
- Raw fruits
- Raw vegetables
- Dried fruit
- Whole grains

Supplies Needed for Procedure Preparation:

- 1) Miralax 238 gram bottle or generic equivalent



- 2) 4 Dulcolax tablets or generic equivalent (not the stool softener or suppositories).

- 3) 64 ounces of electrolyte replacement solution such as Gatorade, Pedialyte, Powerade, or Propel. No red, blue, or purple. You may use sugar free options of these products if you are diabetic.



Split Prep Colonoscopy Procedure Instructions

Day Before Procedure

Step 1:

Start a clear liquid diet. No solid food starting when you wake up the day before your procedure. You may only have clear liquids the day before your procedure. Clear liquids include broth (beef or chicken without any particles), Jello, tea, black coffee (no creamer), Gatorade, Powerade, soda, water. Popsicles, and clear fruit juices (apple, white grape, white cranberry - no pulp). **DO NOT drink milk or milk products. NO red, purple, or blue drinks.**

It is important to stay well hydrated throughout the entire preparation process. Drink plenty of approved clear liquids throughout the day before your procedure and between preparation doses. Good hydration can improve comfort during the prep process and help reduce dehydration-related symptoms such as dizziness, weakness, headaches, cramping, and nausea.

Step 2:

At 3:00 p.m. on the day before your colonoscopy, take 4 Dulcolax tablets all at once with water.

Step 3:

At 5:00 p.m. on the day before your colonoscopy, mix your electrolyte solution preparation. Mix the entire 238-gram bottle of MiraLAX with the 64-ounce electrolyte solution (4-16 oz. bottles). Drink 8 ounces of the prep every 15 minutes until HALF of the prep is gone, then drink two 8-ounce glasses of water.

Step 4:

No hard candies, gum, lozenges, nicotine pouches, chewing tobacco or similar products after midnight.

Important: If you experience prep-related symptoms (examples: nausea, bloating, or cramping), pause or slow down the rate of drinking until your symptoms diminish.

What to Expect During the Preparation

You should begin having frequent bowel movements after starting the preparation. By the time your preparation is complete, your stool should be clear to light yellow liquid with minimal sediment or particles, similar in appearance to urine or clear broth.

If your stool remains brown, thick, or contains significant solid material after completing the preparation, please contact the Endoscopy Department for further instructions, as your preparation may not be adequate for the procedure.

Day of the Procedure



- ### Hydration Reminder

Important: Failure to follow the preparation instructions could result in the procedure being rescheduled or needing to repeat the procedure.

Current Medication List

[illegible]